



## PREVALENCE AND RISK FACTORS OF DRY EYE DISEASE SYMPTOMS AMONG MEDICAL UNDERGRADUATES IN SOUTH INDIA

### Ophthalmology

**Aarudhra Premchander**

Intern Medical Officer, Vydehi Institute of Medical Sciences & RC

**Ananya Chakraborty\***

Professor & HOD, Department of Pharmacology, Vydehi Institute of Medical Sciences & RC \*Corresponding Author

**Vittal Naik**

Professor & HOD, Department of Ophthalmology, Vydehi Institute of Medical Sciences & RC

**Hiren Kolli**

B.S. Virginia Commonwealth University School of Medicine

**Supriya Acharya**

Biostatistician, Department of Community Medicine, Vydehi Institute of Medical Sciences & RC

### ABSTRACT

**BACKGROUND:** The study was conducted to determine the prevalence of symptomatic Dry Eye Disease (DED) its relationship with associated risk factors among medical undergraduates.

**METHODS:** This cross-sectional study was conducted between December 2017 to May 2018. After Institutional Ethics Committee approval, a validated questionnaire was administered to second and third year medical undergraduates. The severity of the disease was measured via the Ocular Surface Disease Index (OSDI) questionnaire.

**RESULTS:** The mean age of the participants ( $\pm$ SD) was 20.33 $\pm$ 1.15 years. The overall prevalence of DED was 39.1%. The prevalence of mild, moderate and severe DED were 21.8% (104), 10% (48) and 7.3% (35) respectively. There was statistically significant association of DED with refractive error, headache, and use of eye drops.

**CONCLUSION:** Symptomatic DED is prevalent amongst medical undergraduates. It is important to diagnose the condition and risk factors early and thereby orient the future doctors towards betterment of their own overall ocular health.

### KEYWORDS

Dry Eye Disease, OSDI, Symptomatic DED

### INTRODUCTION

Dry Eye Disease (DED) is characterized by a vicious cycle of tear film instability, hyperosmolarity, chronic inflammation and neurosensory abnormalities. These subsequently lead to ocular surface damage.<sup>1</sup>

Previous studies have reported several risk factors that influence the development of DED. These are external conditions like low humidity, urban pollution; personal factors like age, systemic diseases; personal activities such excessive use of Video Display Units (VDU), smoking, use of contact lenses, refractive surgery, and use of ocular drops. The reported prevalence of DED ranges from 5 to 50%.<sup>2,3</sup> Clinical signs and symptoms of DED very often lack coherence with subjective symptoms and complaints by the patients and vice versa. The ideal diagnosis of DED, thus, requires a validated symptom questionnaire be administered at the beginning of the patient interaction followed by a battery of clinical tests.<sup>1,4</sup> The Ocular Surface Disease Index Questionnaire (OSDI) is one such questionnaire that is widely used in epidemiological studies.<sup>5</sup> Recent studies conducted using OSDI in medical and other university students from Asian countries stated that symptomatic DED is prevalent in that age group.<sup>2,6,7</sup>

There are many published studies on the ocular health of school children in India. Our literature search in younger population on ocular morbidity in college going students in India yielded very few studies. A recent study from India focused on ocular health in medical and engineering students the context of increased usage of VDU and smart phones.<sup>8</sup> With this background, this study was taken up to determine the prevalence of symptoms of DED and its relationship with associated risk factors in medical students using the OSDI questionnaire.

### METHODS

This cross-sectional study was conducted from December 2017 to May 2018. After the approval from the Institutional ethics committee, undergraduate medical students of Vydehi Institute of Medical Sciences & Research Centre were invited to take part in the study. Permission to use OSDI was obtained from Allergan Inc., Irvine, CA. A 10 item validated questionnaire was developed to estimate the risk factors of DED. The questionnaire along with OSDI was administered

by the authors over 5 sessions. Enough time was given to document responses by the participants. After collecting initial responses, orientation on ocular hygiene was conducted. Participants with OSDI scores suggesting DED were advised to visit Ophthalmology department for further clinical evaluation and needful. The response scores were double checked by the investigators, tabulated, and analyzed using SPSS version 21.

### RESULTS

**Demographic profile:** A total of 500 forms were distributed. 484 forms were returned (response rate: 96.8%). Out of these, 6 forms were incomplete and hence excluded from the study. Finally, a total of 478 responses were evaluated. Out of 478 participants, 68.4% (327) were females, and 31.6% (151) were males. The mean age of the participants ( $\pm$ SD) was 20.33 $\pm$ 1.15 years. The demographic details of participants are shown in Table 1.

**Grouping based on OSDI score:** Based on OSDI score, participants were grouped into normal, mild, moderate, and severe DED respectively. It was found out that 60.9% (291), 21.8% (104), 10% (48) and 7.3% (35) participants had normal, mild, moderate, and severe OSDI scores ( $p < 0.001$ ). The OSDI score based grouping of participants are shown in Table 1.

**Prevalence of DED based on OSDI:** DED was found to be present in 37.6% (123) females and 42.4% (64) males. The association between gender and DED was found to be not significant ( $p$  value was 0.178). The overall prevalence of DED was found to be 39.1%. The prevalence of mild, moderate and severe DED was found to be 21.8% (104), 10% (48) and 7.3% (35) respectively ( $p < 0.001$ ). The results are shown in Table 1.

**Co-morbidities:** Out of all the participants, 64.4% (308) had refractive error. Out of them 52.1% (249) were myopes and 10.5% (50) were hypermetropes. There was a significant difference between OSDI median score (Q1, Q3) in normal, myopes, and hypermetropes respectively ( $p < 0.001$ ). 47.1% (226), 6.3% (30), 47.1 (225), and 5.4% (26) participants suffered from headache, migraine, sinusitis and anemia respectively. It was also found out that 4% (19) participants

were on eye drops. The results are shown in Table 1.

**Table 1: Demographic details, OSDI scores and comorbidities**

Variables	Data	p-value
Age (Mean±SD)	20.33±1.15 years	-
<b>Sex (Percentages/Number)</b>		
Males	31.6% (151)	0.300
Females	68.4 % (327)	
<b>OSDI based groups</b>		
Normal	60.9% (291)	<0.001
Mild DED	21.8% (104)	
Moderate DED	10% (48)	
Severe DED	7.3% (35)	
<b>OSDI median score (Q1, Q3)</b>		
Normal	4.16(2.08,8.33)	<0.001
Mild DED	15(12.77,18.70)	
Moderate DED	27(25,29.01)	
Severe DED	39.2(35.36,50)	
<b>Overall DED sex wise</b>		
Males	42.4% (64)	0.178
Females	37.6% (123)	
<b>Comorbid conditions</b>		
1. Refractive error	64.4% (308)	<0.001
i) Myopia	52.1% (249)	
• Mild (SE between -0.75 and -2.99D)	61.0% (152)	
• Moderate (SE between -0.3.00 and -5.99D)	32.9% (82)	
• Severe (SE equal or more than -6.00)	6.1% (15)	
ii) Hypermetropia	10.5% (50)	
iii) Others	1.8% (9)	
2. Headache	47.1% (225)	
3. Migraine	6.3% (30)	
4. Sinusitis	47.1 (225)	
5. Anemia	5.4% (26)	
<b>OSDI median score (Q1, Q3)</b>		
Normal	5.55(2.03,12.50)	<0.001
Myopes	10.40(4.16,20.45)	
Hypermetropes	9.74(6.25,18.18)	
Use of ophthalmic drops	4% (19)	<0.001

Percentages expressed are out of 478 participants; SD= Standard deviation, Q1: Quartile 1 (25th percentile); Q3: Quartile 3: 75th percentile Kruskal Wallis followed by Mann Whitney U test was performed to compare median OSDI score between the groups. For significant differences in sex, comorbidities, and use of drops Chi Square and Fishers Exact tests were performed where ever appropriate

**Risk factors for symptomatic DED:**

The following risk factors were evaluated. Results are shown in Table 2.

**i) Refractive error:** In participants with refractive error, the overall prevalence of DED was 47% (145). The association of refractive error and DED between the groups was found to be statistically significant (p<0.001).

**ii) Headache:** 47.1% (226) participants mentioned that they suffer from headache, and 6.3 % (30) participants were found to suffer from migraine headache. Among the participants that suffer from headache, the overall prevalence of DED was 48.4% (78). The prevalence of mild, moderate and severe DED was found to be 22.6% (51), 15.1 % (34), and 10.6% (24) respectively. There was statistically significant association of DED with headache (p<0.001).

**iii) Currently on any drops:** 4% (19) participants mentioned that they were on eye drops. Out of them 68.4% (13) were found to have DED. There was statistically significant association of DED with use of eye drops (p<0.001).

**iv) Time spent in reading per day (hours):** 18.4% (88), 27.4% (131), 41.4% (198), 11.9% (57), 0.8% (4) participants were found to spent ≤1 hr, ≤2hr, ≤3hr, ≤4hr, and ≥4hr respectively. The prevalence of DED (overall) was 37.5%, 44.3%, 33.3%, 49.1%, and 50% respectively.

**v) Time spent on TV/Laptop/Smart phone in a day (hours):** 9.8% (47), 26.6 (127), 23.4% (112), 16.3% (78), and 20.5% (98) were found to spent ≤1 hr, ≤2hr, ≤3hr, ≤4hr, and ≥4hr respectively. The prevalence of DED (overall) in the above groups was 31.9%(15), 33.1%(42), 43.8%(49), 50%(39), 39.9%(39) respectively.

**Table 2: Association of risk factors and symptomatic DED**

Questions	Responses	Group					p value
		Normal	Mild	Moderate	Severe	Total	
Do you have refractive error	No	128 (44)	23(22.1)	13(27.1)	6(17.1)	170(35.6)	<0.001
	Yes	163(56)	81(77.9)	35(72.9)	29(82.9)	308(64.4)	
	Total	291(100)	104(100)	48(100)	35(100)	478(100)	
Do you suffer from headache	No	175(60.1)	53(51)	14(29.2)	11(31.4)	253(52.9)	<0.001
	Yes	116(39.9)	51(49)	34(70.8)	24(68.6)	225(47.1)	
	Total	291(100)	104(100)	48(100)	35(100)	478(100)	
Are you currently on any drops	No	285(97.9)	100(96.2)	43(89.6)	31(88.6)	459(96)	0.004
	Yes	6(2.1)	4(3.8)	5(10.4)	4(11.4)	19(4)	
	Total	291(100)	104(100)	48(100)	35(100)	478(100)	
How many hours do you spend in reading in a day	Not Mentioned	14(4.8)	1(1)	1(2.1)	2(5.7)	18(3.8)	NA
	≤1	41(14.1)	17(16.3)	7(14.6)	5(14.3)	70(14.6)	
	≤2	73(25.1)	31(29.8)	17(35.4)	10(28.6)	131(27.4)	
	≤3	132(45.4)	41(39.4)	13(27.1)	12(34.3)	198(41.4)	
	≤4	29(10)	13(12.5)	9(18.8)	6(17.1)	57(11.9)	
	>4	2(0.7)	1(1)	1(2.1)	0(0)	4(0.8)	
	Total	291(100)	104(100)	48(100)	35(100)	478(100)	
How much time do you spend on TV/Laptop/Smart phone in a day(hours)	Not Mentioned	13(4.5)	2(1.9)	0(0)	1(2.9)	16(3.3)	NA
	≤1	32(11)	6(5.8)	5(10.4)	4(11.4)	47(9.8)	
	≤2	85(29.2)	29(27.9)	11(22.9)	2(5.7)	127(26.6)	
	≤3	63(21.6)	29(27.9)	13(27.1)	7(20)	112(23.4)	
	≤4	39(13.4)	19(18.3)	8(16.7)	12(34.3)	78(16.3)	
	>4	59(20.3)	19(18.3)	11(22.9)	9(25.7)	98(20.5)	
	Total	291(100)	104(100)	48(100)	35(100)	478(100)	

Total number of participants: 478; Groups were based on OSDI scores. Statistical analysis was conducted using Chi square test. P value of <0.001 was considered statistically significant.

**DISCUSSION**

This is the first study on symptomatic DED conducted amongst medical undergraduates in India. The prevalence of DED in this study was found to be 39.1%. The prevalence was found to be more in males, but the association was not statistically significant. A study involving 700 university undergraduates in Ghana reported the prevalence of symptomatic DED to be 44.3%. Sex was not significantly associated

with DED.<sup>10</sup> Another cross-sectional study that included 209 students at a medical school in Korea found the prevalence to be 27.1%.<sup>11</sup> A study conducted at a medical school in China estimated the prevalence to be 18.7%. The risk factors evaluated were daily reading time of ≥4 h, daily computer use of ≥4 h and constant use of eyeglasses.<sup>12</sup> A cross-section hospital-based, observational study in North India found out the symptomatic DED prevalence to be 32%. It opined that DED is

more common in the age group of 21–40 years. Male sex (65.3% males, 34.7% females), VDT use of 4 hours or more, and contact lens use were associated with increased odds of developing DED.<sup>13</sup> As our study was on young adults (20.33±1.15 years), the prevalence (39.1%) and male sex preponderance for DED (42.4% males, 36.6% females) coincides with this study conducted in North India.

In this study, the most common refractive error documented was myopia. It was found to be prevalent in 52.1% participants. Previous studies conducted among the medical students of Saudi Arabia, Malaysia, Singapore and Norway also reported higher prevalence of myopia amongst medical students. Our results, thus coincide with the above studies.<sup>14, 15, 16, 17</sup> The Saudi study found out the prevalence of myopia to be 53.7%. Even though genetics has a role to play in myopia, recent studies blame it on lack of ocular hygiene like not taking a break after every hours of studying, studying under a dim lamp, screen time of more than 3 hours/day, rubbing eyes etc. towards increased prevalence of the same.<sup>18</sup> Our study found out that headache, and use of eye drops are risk factors for DED. The results coincide with previous studies.<sup>13, 18, 19</sup>

This study has limitations as follows: 1) Data of only 478 participants from a single medical school were included. This is substantially small number to represent the overall population of medical undergraduates from south India. 2) Closed ended questionnaire was administered to evaluate the risk factors. However, we believe that this study could provide significant information regarding prevalence and risk factors 3) Diagnosis of DED was made based on OSDI, examination for dry eye signs was not performed as part of the study. However, various previous studies were successfully conducted using the questionnaire as the reliability for such is already established. In this study, we suggested to participants to undergo clinical examination. This will help establish the diagnosis, and confirm the severity of DED and plan suitable treatment.

## CONCLUSION

The study provided useful insight about symptomatic DED amongst medical undergraduates from a select medical school of South India. A larger study involving more medical institutions from different geographical regions can be taken up. Based on questionnaire based identification of population at risk for DED, clinical examination can be suggested. This will help prevent ocular morbidities and enhance betterment of overall ocular health.

## REFERENCES

- Craig JP, Nichols KK, Akpek EK, et al. TFOS DEWS II definition and classification report. *Ocul Surf* 2017; 15: pp. 276-283
- Castelyn B, Majola S, Motilal R, et al. Prevalence of dry eye amongst black and Indian university students aged 18–30 years. *African Vision and Eye Health* 2015;74:1-6.
- Rao PD, Kammari P, Shanbhag S et al. Incidence, demographics, types and risk factors of dry eye disease in India: Electronic medical records driven big data analytics report I. *Ocular Surface* 2019; 17(2): 250-256.
- Amparo F, Schaumberg DA, Dana R. Comparison of Two Questionnaires for Dry Eye Symptom Assessment: The ocular surface disease index and the symptom assessment in dry eye. *Ophthalmology* 2015; 122(7):1498–1503.
- Schiffman RM, Christianson MD, Jacobsen G, et al. Reliability and validity of the ocular surface disease index. *Arch Ophthalmol* 2000; 118:615–621.
- Hyon JY, Yang HK, Han SB. Dry Eye Symptoms May Have Association With Psychological Stress in Medical Students. *Eye Contact Lens* 2019;45(5):310-314.
- Asiedu K, Kvei S, Boampong F, et al. Symptomatic dry eye and its associated factors: A study of university undergraduate students in Ghana. *Eye Contact Lens* 2017 Jul;43(4):262-266.
- Gupta M, Gupta BP, Chauhan A, et al. Ocular morbidity prevalence among school children in Shimla, Himachal, North India. *Indian J Ophthalmol*. 2009;57(2):133–138.
- Logaraj M, Madhupriya V, Hegde S. Computer vision syndrome and associated factors among medical and engineering students in Chennai. *Ann Med Health Sci Res*. 2014;4(2):179–185.
- Asiedu K, Kyei S, Boampong F, et al., Symptomatic dry eye and its associated factors: A study of University undergraduate students in Ghana. *Eye Contact Lens* 2017 Jul;43(4):262-266.
- Hyon JY, Yang HK, Han SB. Dry eye symptoms may have association with psychological stress in medical students. *Eye Contact Lens* 2019 Sep;45(5):310-314.
- He YP, Zhang WF, L P, Zhou R, Xia JT, Fan Y. Prevalence of symptomatic dry eye disease among Chinese college students with associated risk factors. *Int Eye Sci* 2016;16(6):1019-1025.
- Singh TJ, Clara FR, Kaur M, et al., Prevalence and risk factors of dry eye disease in North India: Ocular surface disease index-based cross-sectional hospital study. *Ind J Ophthal* 2018; 66(2): 207-211.
- Al-Rashidi SH, Albahouth AA, Althwini WA, et al. Prevalence of refractive errors among medical students of Qassim University, Saudi Arabia: Cross-sectional descriptive study. *Open Access Maced J Med Sci*. 2018;6(5):940–943.
- Gopalakrishnan S, Prakash MVS, Jha K, Ranjit A study of refractive errors among medical students in AIMST University, Malaysia. *Indian Medical Journal* 2011; 105:365–367.
- Midelfart A, Aamo B, Sjøhaug KA, Dysthe BE. Myopia among medical students in Norway. *Acta Ophthalmol* 1992;70:317–22.
- Chow YC, Dhillon B, Chew PTK, Chew SJ. Refractive errors in Singapore medical students. *Singapore Med J* 1990;31:472–3.
- Wang, L., Du, M., Yi, H. et al. Prevalence of and Factors Associated with Myopia in

Inner Mongolia Medical Students in China, a cross-sectional study. *BMC Ophthalmol* 17, 52 (2017).

- Koktekir BE, Celik G, Karalezli A. et al., Dry eyes and migraines: is there really a correlation? *Cornea* 2012;31(12):1414-6.